

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
5 December 2002 (05.12.2002)

PCT

(10) International Publication Number  
**WO 02/096885 A1**

(51) International Patent Classification<sup>7</sup>: **C07D 239/34**,  
401/12, A61K 31/505, A61P 29/00

(21) International Application Number: PCT/GB02/02415

(22) International Filing Date: 23 May 2002 (23.05.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
0112802.4 25 May 2001 (25.05.2001) GB

(71) Applicant (for all designated States except US): **GLAXO GROUP LIMITED** [GB/GB]; Glaxo Wellcome House, Berkeley Avenue, Greenford, Middlesex UB6 ONN (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **NAYLOR, Alan**

[GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage, Hertfordshire SG1 2NY (GB). **PAYNE, Jeremy, John** [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage, Hertfordshire SG1 2NY (GB). **PEGG, Neil, Anthony** [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage, Hertfordshire SG1 2NY (GB).


(74) Agent: **GIDDINGS, Peter, John**; Corporate Intellectual Property, GlaxoSmithKline, 980 Great West Road (CN925.1), Brentford, Middlesex TW8 9GS (GB).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),

[Continued on next page]

(54) Title: PYRIMIDINE DERIVATIVES USEFUL AS SELECTIVE COX-2 INHIBITORS



WO 02/096885 A1

(57) Abstract: The invention provides the compounds of formula (I) in which: R<sup>1</sup> is selected from the group consisting of H, C<sub>1-6</sub>alkyl, C<sub>1-2</sub>alkyl substituted by one to five fluorine atoms, C<sub>3-6</sub>alkenyl, C<sub>3-10</sub>cycloalkyl, C<sub>0-6</sub>alkyl, C<sub>4-12</sub>bridged cycloalkyl, A(CR<sup>4</sup>R<sup>5</sup>)<sub>n</sub> and b(CR<sup>4</sup>R<sup>5</sup>)<sub>n</sub>; R<sup>2</sup> is C<sub>1-2</sub>alkyl substituted by one to five fluorine atoms; R<sup>3</sup> is selected from the group consisting of C<sub>1-6</sub>alkyl, NH<sub>2</sub> and R<sup>7</sup>CONH; R<sup>4</sup> and R<sup>5</sup> are independently selected from H or C<sub>1-6</sub>alkyl; A is an unsubstituted 5- or 6-membered heteroaryl or an unsubstituted 6-membered aryl, or a 5- or 6-membered heteroaryl or a 6-membered aryl substituted by one or more R<sup>6</sup>; R<sup>6</sup> is selected from the group consisting of halogen, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyl substituted by one or more fluorine atoms, C<sub>1-6</sub>alkoxy, C<sub>1-6</sub>alkoxy substituted by one or more F, NH<sub>2</sub>SO<sub>2</sub> and C<sub>1-6</sub>alkylSO<sub>2</sub>; B is selected from the group consisting of Formula (i) and (ii) and where (iv) defines the point of attachment of the ring; R<sup>7</sup> is selected from the group consisting of H, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkoxy, C<sub>1-6</sub>alkylOC<sub>1-6</sub>alkyl, phenyl, HO<sub>2</sub>CC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkylOCOC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkylOCO, H<sub>2</sub>NC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkylOCONHC<sub>1-6</sub>alkyl and C<sub>1-6</sub>alkylCONHC<sub>1-6</sub>alkyl; and n is 0 to 4. Compounds of formula (I) are potent and selective inhibitors of COX-2 and are of use in treatment of the pain, fever and inflammation of variety of conditions and diseases.



Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,  
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent  
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,  
NE, SN, TD, TG).

— *before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendments*

**Published:**

— *with international search report*


*For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.*

## PYRIMIDINE DERIVATIVES USEFUL AS SELECTIVE COX-2 INHIBITORS

This invention relates to pyrimidine derivatives, to processes for their preparation, to pharmaceutical compositions containing them and to their use in medicine.

5 The enzyme cyclooxygenase (COX) has recently been discovered to exist in two isoforms, COX-1 and COX-2. COX-1 corresponds to the originally identified constitutive enzyme while COX-2 is rapidly and readily inducible by a number of agents including mitogens, endotoxin, hormones, cytokines and growth factors. Prostaglandins generated by the action of COX have both physiological and pathological roles. It is generally believed that COX-1 is largely responsible for the important physiological functions such as maintenance of gastrointestinal integrity and renal blood flow. In contrast the inducible form, COX-2, is believed to be largely responsible for the pathological effects of prostaglandins where rapid induction of the enzyme occurs in response to such agents as inflammatory agents, hormones, growth factors and cytokines. A selective inhibitor of COX-2 would therefore have anti-inflammatory, anti-pyretic and analgesic properties, without the potential side effects associated with inhibition of COX-1. We have now found a novel group of compounds which are both potent and selective inhibitors of COX-2.

20 The invention thus provides the compounds of formula (I)



in which:

25 R¹ is selected from the group consisting of H, C<sub>1-6</sub>alkyl, C<sub>1-2</sub>alkyl substituted by one to five fluorine atoms, C<sub>3-6</sub>alkenyl, C<sub>3-6</sub>alkynyl, C<sub>3-10</sub>cycloalkyl/C<sub>0-6</sub>alkyl, C<sub>4-12</sub>bridged cycloalkyl, A(CR⁴R⁵)<sub>n</sub> and B(CR⁴R⁵)<sub>n</sub>;

R² is C<sub>1-2</sub>alkyl substituted by one to five fluorine atoms;


R³ is selected from the group consisting of C<sub>1-6</sub>alkyl, NH<sub>2</sub> and R⁷CONH;

$R^4$  and  $R^5$  are independently selected from H or  $C_{1-6}$ alkyl;

A is an unsubstituted 5- or 6-membered heteroaryl or an unsubstituted 6-membered aryl, or a 5- or 6-membered heteroaryl or a 6-membered aryl substituted by one or more  $R^6$ ;

- 5  $R^6$  is selected from the group consisting of halogen,  $C_{1-6}$ alkyl,  $C_{1-6}$ alkyl substituted by one more fluorine atoms,  $C_{1-6}$ alkoxy,  $C_{1-6}$ alkoxy substituted by one or more F,  $NH_2SO_2$  and  $C_{1-6}alkylSO_2$ ;

B is selected from the group consisting of



Y defines the point of attachment of the ring;


- 10  $R^7$  is selected from the group consisting of H,  $C_{1-6}$ alkyl,  $C_{1-6}$ alkoxy,  $C_{1-6}alkylOC_{1-6}alkyl$ , phenyl,  $HO_2CC_{1-6}alkyl$ ,  $C_{1-6}alkylOCOC_{1-6}alkyl$ ,  $C_{1-6}alkylOCO$ ,  $H_2NC_{1-6}alkyl$ ,  $C_{1-6}alkylOCONHC_{1-6}alkyl$  and  $C_{1-6}alkylCONHC_{1-6}alkyl$ ; and n is 0 to 4.

The term halogen is used to represent fluorine, chlorine, bromine or iodine.


- 15 The term 'alkyl' as a group or part of a group means a straight or branched chain alkyl group, for example a methyl, ethyl, n-propyl, i-propyl, n-butyl, s-butyl or t-butyl group.

The term 5-membered heteroaryl means a heteroaryl selected from the following:


20



The term 6- membered heteroaryl means a heteroaryl selected from the following:



5 The term 6-membered aryl means:



10 It is to be understood that the present invention encompasses all isomers of the compounds of formula (I) and their pharmaceutically acceptable derivatives, including all geometric, tautomeric and optical forms, and mixtures thereof (e.g. racemic mixtures). In particular when the ring B lacks a plane of symmetry the compounds of formula (I) contain a chiral centre as indicated therein by the asterisk \*. Furthermore, it will be appreciated by those skilled in the art that when  $R^4$  and  $R^5$  in formula (I) are different the corresponding compounds contain at least one chiral centre, by virtue of the asymmetric carbon atom defined thereby, and that such compounds exist in the form of a pair of optical isomers (i.e. enantiomers).

In one aspect of the invention  $R^1$  is selected from the group consisting of H,  $C_{1-6}$ alkyl,  $C_{1-2}$ alkyl substituted by one to five fluorine atoms,  $C_{3-6}$ alkenyl,  $C_{3-6}$ alkynyl,  $C_{3-10}$ cycloalkyl,  $C_{0-6}$ alkyl,  $C_{4-12}$ bridged cycloalkyl and  $B(CR^4R^5)_n$ ;

20 In another aspect of the invention  $R^1$  is  $C_{1-6}$ alkyl or  $C_{1-2}$ alkyl substituted by one to five fluorine atoms. In another aspect  $R^1$  is  $C_{2-6}$ alkyl (e.g. n-butyl).

In another aspect of the invention  $R^1$  is  $C_{3-10}$ cycloalkyl, such as  $C_{3-10}$ cycloalkyl (e.g. cyclopentyl or cyclohexyl). In another aspect  $R^1$  is  $C_{3-10}$ cycloalkylmethyl, such as  $C_{3-7}$ cycloalkylmethyl (e.g. cyclopentylmethyl).


25 In another aspect of the invention  $R^1$  is  $A(CR^4R^5)_n$ .

In another aspect of the invention  $R^2$  is  $\text{CHF}_2$ ,  $\text{CH}_2\text{F}$  or  $\text{CF}_3$ . In another aspect  $R^2$  is  $\text{CF}_3$ .

In another aspect of the invention  $R^3$  is  $\text{C}_{1-6}$ alkyl, such as  $\text{C}_{1-3}$ alkyl (e.g. methyl).

5 In another aspect of the invention  $R^4$  and  $R^5$  are independently selected from H or methyl. In another aspect  $R^4$  and  $R^5$  are both H.

In another aspect of the invention A is selected from the group consisting of



) defines the point of attachment of the ring  
and A is unsubstituted or substituted by one or two  $R^6$ .

10 In another aspect of the invention  $R^6$  is selected from the group consisting of halogen (e.g. F),  $\text{C}_{1-3}$ alkyl (e.g. methyl),  $\text{C}_{1-3}$ alkyl substituted by one to three fluorine atoms (e.g.  $\text{CF}_3$ ), and  $\text{C}_{1-3}$ alkoxy (e.g. methoxy).

In another aspect of the invention  $R^7$  is selected from the group consisting of  $\text{C}_{1-6}$ alkyl (e.g. ethyl), phenyl and aminomethyl.

In another aspect of the invention n is 1 to 4.

15 In another aspect of the invention n is 0 to 2 (e.g. 0).


It is to be understood that the invention covers all combinations of particular aspects of the invention as described hereinabove.

20 Within the invention there is provided one group of compounds of formula (I) (group A) wherein:  $R^1$  is  $\text{C}_{1-6}$ alkyl (e.g. n-butyl);  $R^2$  is  $\text{CF}_3$ ; and  $R^3$  is  $\text{C}_{1-6}$ alkyl, such as  $\text{C}_{1-3}$ alkyl (e.g. methyl).

Within the invention there is provided another group of compounds of formula (I) (group B) wherein:  $R^1$  is  $\text{C}_{3-10}$ cycloalkyl $\text{C}_{0-6}$ alkyl, such as  $\text{C}_{3-10}$ cycloalkyl (e.g. cyclopentyl or cyclohexyl);  $R^2$  is  $\text{CF}_3$ ; and  $R^3$  is  $\text{C}_{1-6}$ alkyl, such as  $\text{C}_{1-3}$ alkyl (e.g. methyl).


Within the invention there is provided another group of compounds of formula (I) (group C) wherein:  $R^1$  is  $C_{3-10}$ cycloalkylmethyl, such as  $C_{3-7}$ cycloalkylmethyl (e.g. cyclopentylmethyl);  $R^2$  is  $CF_3$ ; and  $R^3$  is  $C_{1-6}$ alkyl, such as  $C_{1-3}$ alkyl (e.g. methyl).

5 Within the invention there is provided another group of compounds of formula (I) (group D) wherein:  $R^1$  is  $A(CR^4R^5)_n$ ;  $R^2$  is  $CF_3$ ;  $R^3$  is  $C_{1-6}$ alkyl, such as  $C_{1-3}$ alkyl (e.g. methyl);  $R^4$  and  $R^5$  are independently selected from H or methyl; A is selected from the group consisting of



10 and A is unsubstituted or substituted by one or two  $R^6$ ;  $R^6$  is selected from the group consisting of halogen (e.g. F),  $C_{1-3}$ alkyl (e.g. methyl),  $C_{1-3}$ alkyl substituted by one to three fluorine atoms (e.g.  $CF_3$ ), and  $C_{1-3}$ alkoxy (e.g. methoxy); and n is 0 to 2 (e.g. 0).

15 Within group D, there is provided a further group of compounds (group D1) wherein:  $R^1$  is  $A(CR^4R^5)_n$ ;  $R^2$  is  $CF_3$ ;  $R^3$  is methyl;  $R^4$  and  $R^5$  are both H; A is selected from the group consisting of



and A is unsubstituted or substituted by one or two  $R^6$ ;  $R^6$  is selected from the group consisting of fluorine, chlorine, methyl,  $CF_3$  and methoxy; and n is 0 or 1.

In a preferred aspect the invention provides the following compounds:

- 20 2-(4-fluorophenoxy)-4-[4-(methylsulfonyl)phenyl]-6-(trifluoromethyl)pyrimidine;  
 2-(4-methoxyphenoxy)-4-[4-(methylsulfonyl)phenyl]-6-trifluoromethyl)pyrimidine;  
 2-butoxy-4-[4-(methylsulfonyl)phenyl]-6-(trifluoromethyl)pyrimidine;  
 2-[(5-chloropyridin-3-yl)oxy]-4-[4-(methylsulfonyl)phenyl]-6-  
 (trifluoromethyl)pyrimidine;  
 25 2-(cyclohexyloxy)-4-[4-(methylsulfonyl)phenyl]-6-(trifluoromethyl)pyrimidine.

In a more preferred aspect the invention provides the following compound:

2-butoxy-4-[4-(methylsulfonyl)phenyl]-6-(trifluoromethyl)pyrimidine.

5 Since the compounds of the present invention, in particular compounds of formula (I), are intended for use in pharmaceutical compositions, it will be understood that they are each provided in substantially pure form, for example at least 50% pure, more suitably at least 75% pure and preferably at least 95% pure (% are on a wt/wt basis). Impure preparations of the compound of formula (I) may be used for preparing the more pure forms used in pharmaceutical compositions. Although the purity of intermediate compounds of the present  
10 invention is less critical, it will be readily understood that the substantially pure form is preferred as for the compounds of formula (I). Preferably, whenever possible, the compounds of the present invention are available in crystalline form.

15 When some of the compounds of this invention are allowed to crystallise or are recrystallised from organic solvents, solvent of recrystallisation may be present in the crystalline product. This invention includes within its scope such solvates. Similarly, some of the compounds of this invention may be crystallised or recrystallised from solvents containing water. In such cases water of hydration may be formed. This invention includes within its scope stoichiometric hydrates  
20 as well as compounds containing variable amounts of water that may be produced by processes such as lyophilisation. In addition, different crystallisation conditions may lead to the formation of different polymorphic forms of crystalline products. This invention includes within its scope all the polymorphic forms of the compounds of formula (I).

25 Compounds of the invention are potent and selective inhibitors of COX-2. This activity is illustrated by their ability to selectively inhibit COX-2 over COX-1.

In view of their selective COX-2 inhibitory activity, the compounds of the present invention are of interest for use in human and veterinary medicine, particularly in the treatment of the pain (both chronic and acute), fever and inflammation of a  
30 variety of conditions and diseases mediated by selective inhibition of COX-2. Such conditions and diseases are well known in the art and include rheumatic fever; symptoms associated with influenza or other viral infections, such as the common cold; lower back and neck pain; headache; toothache; sprains and strains; myositis; sympathetically maintained pain; synovitis; arthritis, including

rheumatoid arthritis; degenerative joint diseases, including osteoarthritis; gout and ankylosing spondylitis; tendinitis; bursitis; skin related conditions, such as psoriasis, eczema, burns and dermatitis; injuries, such as sports injuries and those arising from surgical and dental procedures.

5 The compounds of the invention are also useful for the treatment of neuropathic pain. Neuropathic pain syndromes can develop following neuronal injury and the resulting pain may persist for months or years, even after the original injury has healed. Neuronal injury may occur in the peripheral nerves, dorsal roots, spinal  
10 cord or certain regions in the brain. Neuropathic pain syndromes are traditionally classified according to the disease or event that precipitated them. Neuropathic pain syndromes include: diabetic neuropathy; sciatica; non-specific lower back pain; multiple sclerosis pain; fibromyalgia; HIV-related neuropathy; neuralgia, such as post-herpetic neuralgia and trigeminal neuralgia; and pain resulting from physical trauma, amputation, cancer, toxins or chronic  
15 inflammatory conditions. These conditions are difficult to treat and although several drugs are known to have limited efficacy, complete pain control is rarely achieved. The symptoms of neuropathic pain are incredibly heterogeneous and are often described as spontaneous shooting and lancinating pain, or ongoing, burning pain. In addition, there is pain associated with normally non-painful  
20 sensations such as "pins and needles" (paraesthesias and dysesthesias), increased sensitivity to touch (hyperesthesia), painful sensation following innocuous stimulation (dynamic, static or thermal allodynia), increased sensitivity to noxious stimuli (thermal, cold, mechanical hyperalgesia), continuing pain sensation after removal of the stimulation (hyperpathia) or an absence of or  
25 deficit in selective sensory pathways (hypoalgesia).

The compounds of the invention are also useful for the treatment of other conditions mediated by selective inhibition of COX-2.

For example, the compounds of the invention inhibit cellular and neoplastic transformation and metastatic tumour growth and hence are useful in the  
30 treatment of certain cancerous diseases, such as colonic cancer and prostate cancer. The compounds of the invention are also useful in reducing the number of adenomatous colorectal polyps and thus reduce the risk of developing colon cancer. The compounds of the invention are also useful in the treatment of

cancer associated with overexpression of HER-2/neu, in particular breast cancer.

5 Compounds of the invention also prevent neuronal injury by inhibiting the generation of neuronal free radicals (and hence oxidative stress) and therefore are of use in the treatment of stroke; epilepsy; and epileptic seizures (including grand mal, petit mal, myoclonic epilepsy and partial seizures).

Compounds of the invention also inhibit prostanoid-induced smooth muscle contraction and hence are of use in the treatment of dysmenorrhoea and premature labour.

10 Compounds of the invention are also useful in the treatment of liver disease, such as inflammatory liver disease, for example chronic viral hepatitis B, chronic viral hepatitis C, alcoholic liver injury, primary biliary cirrhosis, autoimmune hepatitis, nonalcoholic steatohepatitis and liver transplant rejection.

15 Compounds of the invention inhibit inflammatory processes and therefore are of use in the treatment of asthma, allergic rhinitis and respiratory distress syndrome; gastrointestinal conditions such as inflammatory bowel disease, Crohn's disease, gastritis, irritable bowel syndrome and ulcerative colitis; and the inflammation in such diseases as vascular disease, migraine, periarteritis nodosa, thyroiditis, aplastic anaemia, Hodgkin's disease, sclerodoma, type I  
20 diabetes, myasthenia gravis, multiple sclerosis, sarcoidosis, nephrotic syndrome, Bechet's syndrome, polymyositis, gingivitis, conjunctivitis and myocardial ischemia.

25 Compounds of the invention are also useful in the treatment of ophthalmic diseases such as retinitis, retinopathies, uveitis and of acute injury to the eye tissue.

30 Compounds of the invention are also useful for the treatment of cognitive disorders such as dementia, particularly degenerative dementia (including senile dementia, Alzheimer's disease, Pick's disease, Huntington's chorea, Parkinson's disease and Creutzfeldt-Jakob disease), and vascular dementia (including multi-infarct dementia), as well as dementia associated with intracranial space occupying lesions, trauma, infections and related conditions (including HIV infection), metabolism, toxins, anoxia and vitamin deficiency; and mild cognitive

impairment associated with ageing, particularly Age Associated Memory Impairment.

5 Compounds of the invention are also useful in the treatment of disorders ameliorated by a gastroprokinetic agent. Disorders ameliorated by gastroprokinetic agents include ileus, for example post-operative ileus and ileus during sepsis; gastroesophageal reflux disease (GORD, or its synonym GERD); gastroparesis, such as diabetic gastroparesis; and other functional bowel disorders, such as non-ulcerative dyspepsia (NUD) and non-cardiac chest pain (NCCP).

10 According to a further aspect of the invention, we provide a compound of formula (I) for use in human or veterinary medicine.

According to another aspect of the invention, we provide a compound of formula (I) for use in the treatment of a condition which is mediated by COX-2.

15 According to a further aspect of the invention, we provide a method of treating a human or animal subject suffering from a condition which is mediated by COX-2 which comprises administering to said subject an effective amount of a compound of formula (I).

20 According to a further aspect of the invention, we provide a method of treating a human or animal subject suffering from an inflammatory disorder, which method comprises administering to said subject an effective amount of a compound of formula (I).

According to another aspect of the invention, we provide the use of a compound of formula (I) for the manufacture of a therapeutic agent for the treatment of a condition which is mediated by COX-2.

25 According to another aspect of the invention, we provide the use of a compound of formula (I) for the manufacture of a therapeutic agent for the treatment of an inflammatory disorder.

30 It is to be understood that reference to treatment includes both treatment of established symptoms and prophylactic treatment, unless explicitly stated otherwise.

It will be appreciated that the compounds of the invention may advantageously be used in conjunction with one or more other therapeutic agents. Examples of suitable agents for adjunctive therapy include a 5HT<sub>1</sub> agonist, such as a triptan (e.g. sumatriptan or naratriptan); an adenosine A<sub>1</sub> agonist; an EP ligand; an NMDA modulator, such as a glycine antagonist; a sodium channel blocker (e.g. lamotrigine); a substance P antagonist (e.g. an NK<sub>1</sub> antagonist); a cannabinoid; acetaminophen or phenacetin; a 5-lipoxygenase inhibitor; a leukotriene receptor antagonist; a DMARD (e.g. methotrexate); gabapentin and related compounds; a tricyclic antidepressant (e.g. amitriptyline); a neurone stabilising antiepileptic drug; a mono-aminergic uptake inhibitor (e.g. venlafaxine); a matrix metalloproteinase inhibitor; a nitric oxide synthase (NOS) inhibitor, such as an iNOS or an nNOS inhibitor; an inhibitor of the release, or action, of tumour necrosis factor  $\alpha$ ; an antibody therapy, such as a monoclonal antibody therapy; an antiviral agent, such as a nucleoside inhibitor (e.g. lamivudine) or an immune system modulator (e.g. interferon); an opioid analgesic; a local anaesthetic; a stimulant, including caffeine; an H<sub>2</sub>-antagonist (e.g. ranitidine); a proton pump inhibitor (e.g. omeprazole); an antacid (e.g. aluminium or magnesium hydroxide); an antifatulent (e.g. simethicone); a decongestant (e.g. phenylephrine, phenylpropanolamine, pseudoephedrine, oxymetazoline, epinephrine, naphazoline, xylometazoline, propylhexedrine, or levo-desoxyephedrine); an antitussive (e.g. codeine, hydrocodone, carmiphen, carbetapentane, or dextromethorphan); a diuretic; or a sedating or non-sedating antihistamine. It is to be understood that the present invention covers the use of a compound of formula (I) in combination with one or more other therapeutic agents.

The compounds of formula (I) are conveniently administered in the form of pharmaceutical compositions. Thus, in another aspect of the invention, we provide a pharmaceutical composition comprising a compound of formula (I) adapted for use in human or veterinary medicine. Such compositions may conveniently be presented for use in conventional manner in admixture with one or more physiologically acceptable carriers or excipients.

The compounds of formula (I) may be formulated for administration in any suitable manner. They may, for example, be formulated for topical administration or administration by inhalation or, more preferably, for oral, transdermal or parenteral administration. The pharmaceutical composition may

be in a form such that it can effect controlled release of the compounds of formula (I).

5 For oral administration, the pharmaceutical composition may take the form of, for example, tablets (including sub-lingual tablets), capsules, powders, solutions, syrups or suspensions prepared by conventional means with acceptable excipients.

For transdermal administration, the pharmaceutical composition may be given in the form of a transdermal patch, such as a transdermal iontophoretic patch.

10 For parenteral administration, the pharmaceutical composition may be given as an injection or a continuous infusion (e.g. intravenously, intravascularly or subcutaneously). The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles and may contain formulatory agents such as suspending, stabilising and/or dispersing agents. For administration by injection these may take the form of a unit dose presentation or  
15 as a multidose presentation preferably with an added preservative.

Alternatively for parenteral administration the active ingredient may be in powder form for reconstitution with a suitable vehicle.

20 The compounds of the invention may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds of the invention may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

25 As stated above, the compounds of the invention may also be used in combination with other therapeutic agents. The invention thus provides, in a further aspect, a combination comprising a compound of formula (I) together with a further therapeutic agent.

30 The combinations referred to above may conveniently be presented for use in the form of a pharmaceutical formulation and thus pharmaceutical formulations comprising a combination as defined above together with a pharmaceutically acceptable carrier or excipient comprise a further aspect of the invention. The

individual components of such combinations may be administered either sequentially or simultaneously in separate or combined pharmaceutical formulations.


5 When a compound of formula (I) is used in combination with a second therapeutic agent active against the same disease state the dose of each compound may differ from that when the compound is used alone. Appropriate doses will be readily appreciated by those skilled in the art.

10 A proposed daily dosage of a compound of formula (I) for the treatment of man is 0.01mg/kg to 500mg/kg, such as 0.05mg/kg to 100mg/kg, e.g. 0.1mg/kg to 50mg/kg, which may be conveniently administered in 1 to 4 doses. The precise dose employed will depend on the age and condition of the patient and on the route of administration. Thus, for example, a daily dose of 0.25mg/kg to 10mg/kg may be suitable for systemic administration.

15 Compounds of formula (I) may be prepared by any method known in the art for the preparation of compounds of analogous structure.

Compounds of formula (I) may be prepared by a process which comprises:

reacting an alcohol  $R^1OH$  of formula (II) or a protected derivative thereof with a compound of formula (III)



20 and thereafter and if necessary,  
interconverting a compound of formula (I) into another compound of formula (I);  
and/or

deprotecting a protected derivative of compound of formula (I).

25 The overall synthesis of a compound of formula (I) is shown in Scheme 1 below in which,  $R^1$  and  $R^2$  are as defined in formula (I) above unless otherwise stated,

$R^3$  is  $C_{1-6}$ alkyl; THF is tetrahydrofuran; MTBE is methyl t-butyl ether; and alkyl is a straight or branched chain alkyl group, for example a methyl, ethyl, n-propyl, i-propyl, n-butyl, s-butyl or t-butyl group.

5 Referring to Scheme 1, the preparation of compounds of formula (I) may conveniently be achieved by the treatment of compounds of formula (III) with an alcohol of formula (II) in the presence of sodium hydride. The reaction is conveniently carried out in a solvent such as THF and at between ambient temperature and reflux.

10 Conveniently the oxidation shown in Scheme 1 is effected using a monopersulfate compound, such as potassium peroxymonosulfate (known as Oxone<sup>TM</sup>) and the reaction is carried out in a solvent, such as an aqueous alcohol, (e.g. aqueous methanol), and at between -78°C and ambient temperature.


15 Alternatively, the oxidation shown in Scheme 1 may be effected using hydrogen peroxide in the presence of catalytic sodium tungstate dihydrate. The reaction may be carried out in a solvent such as acetic acid and at between ambient temperature and reflux (e.g. 50°C).

20 Referring to Scheme 1, the cyclisation of diones of formula (VI) to give the corresponding pyrimidines of formula (IV) is conveniently carried out employing a thioronium salt such as a 2-methyl-2-thiopseudourea sulfate and under reflux.

It will be appreciated by those skilled in the art that certain of the procedures described in Scheme 1 for the preparation of compounds of formula (I) or intermediates thereto may not be applicable to some of the possible substituents.


25 It will be further appreciated by those skilled in the art that it may be necessary or desirable to carry out the transformations described in Scheme 1 in a different order from that described, or to modify one or more of the transformations, to provide the desired compound of formula (I).

Scheme 1



- 5 In one variation of Scheme 1, compounds of formula (III) wherein  $\text{R}^3$  is  $\text{C}_{1-6}\text{alkyl}$  or  $\text{NH}_2$  may be prepared by oxidising a compound of formula (IV)A:

15



under oxidation conditions described hereinabove. Compounds of formula (IV)A may be prepared according to the general procedures of Scheme 1 by employing sulphonyl derivatives in place of the corresponding sulfide compounds of formulae (VI) and (VII).

It will be appreciated by those skilled in the art that compounds of formula (I) may be prepared by interconversion, utilising other compounds of formula (I) as precursors. Suitable interconversions, such as alkylations, are well known to those skilled in the art and are described in many standard organic chemistry texts, such as 'Advanced Organic Chemistry' by Jerry March, fourth edition (Wiley, 1992), incorporated herein by reference. For example, compounds of formula (I) wherein R<sup>1</sup> is C<sub>1-6</sub>alkyl, C<sub>1-2</sub>alkyl substituted by one to five fluorine atoms, C<sub>3-6</sub>alkenyl, C<sub>3-6</sub>alkynyl, C<sub>3-10</sub>cycloalkyl, C<sub>0-6</sub>alkyl, C<sub>4-12</sub>bridged cycloalkane, A(CR<sup>4</sup>R<sup>5</sup>)<sub>n</sub> (with the proviso that n is not zero) and B(CR<sup>4</sup>R<sup>5</sup>)<sub>n</sub> may be prepared by alkylating the corresponding compound of formula (I) wherein R<sup>1</sup> is H.

Acylation of compounds of formula (I) wherein R<sup>3</sup> is NH<sub>2</sub>, to provide compounds of formula (I) wherein R<sup>3</sup> is R<sup>7</sup>CONH, may be carried out by conventional means, for example by employing conventional acylating agents such as those described in 'Advanced Organic Chemistry', pp 417-424, incorporated herein by reference.

As will be appreciated by those skilled in the art it may be necessary or desirable at any stage in the synthesis of compounds of formula (I) to protect one or more sensitive groups in the molecule so as to prevent undesirable side reactions. The protecting groups used in the preparation of compounds of formula (I) may be used in conventional manner. See, for example, those described in 'Protective Groups in Organic Synthesis' by Theodora W Green and Peter G M Wuts, second edition, (John Wiley and Sons, 1991), incorporated herein by reference, which also describes methods for the removal of such groups.

Alcohols of formula (II) are either known compounds or may be prepared by literature methods, such as those described in 'Comprehensive Organic Transformations: a guide to functional group preparations' by Richard Larock (VCH, 1989), incorporated herein by reference.

- 5 Thioronium salts of formula (V) are either known compounds or may be prepared by literature methods, such as those described in A H Owens *et al*, Eur J Med Chem, 1988, 23(3), 295-300, incorporated herein by reference.

Acetophenones of formula (VII) are either known compounds or may be prepared by conventional chemistry.

- 10 Certain intermediates described above are novel compounds, and it is to be understood that all novel intermediates herein form further aspects of the present invention. Compounds of formulae (III) and (IV) are key intermediates and represent a particular aspect of the present invention.

- 15 Solvates (e.g. hydrates) of a compound of the invention may be formed during the work-up procedure of one of the aforementioned process steps.

- The Intermediates and Examples that follow illustrate the invention but do not limit the invention in any way. All temperatures are in °C. Flash column chromatography was carried out using Merck 9385 silica. Solid Phase Extraction (SPE) chromatography was carried out using Varian Mega Bond Elut (Si) cartridges (Anachem) under 15mmHg vacuum. Thin layer chromatography (Tlc) was carried out on silica plates. In addition to those already defined, the following abbreviations are used: Me, methyl; Ac, acyl; DMSO, dimethylsulphoxide; TFA, trifluoroacetic acid; DME, dimethoxyethane; DCM, dichloromethane; NMP, N- methyl pyrrolidone; and MTBE, methyl t-butyl ether.
- 20
- 25

Intermediate 14,4,4-Trifluoro-1-[4-(methylthio)phenyl]butane-1,3-dione

To a solution of ethyl trifluoroacetate (7.95ml, 1.1eq) in MTBE (125ml) was added dropwise 25% sodium methoxide in methanol (16ml, 1.2eq). 4-Methylthioacetophenone (Aldrich, 10g, 0.06mol) was added portionwise and the mixture stirred at ambient temperature overnight. 2N Hydrochloric acid (40ml) was added cautiously and the organic phase separated, washed with brine and dried (Na<sub>2</sub>SO<sub>4</sub>) to give an orange solid. The orange solid was recrystallised from hot isopropanol to give the title compound as a yellow crystalline solid (11.25g, 71%).

MH- 261

Intermediate 22-(Methylthio)-4-[4-(methylthio)phenyl]-6-(trifluoromethyl) pyrimidine

To a mixture of 4,4,4-trifluoro-1-[4-(methylthio)phenyl]butane-1,3-dione (5g) and 2-methyl-2-thiopseudourea sulfate (5.1g, 0.98eq) in acetic acid (100ml) was added sodium acetate (3g, 2eq) and heated under reflux for 8h. The mixture was concentrated *in vacuo* and water (100ml) added to give a solid, which was isolated by filtration to give the title compound as a yellow solid (5.8g, quantitative).

MH+ 317

Intermediate 32-(Methylsulfonyl)-4-[4-(methylsulfonyl)phenyl]-6-(trifluoromethyl)pyrimidine

To a solution of 2-(methylthio)-4-[4-(methylthio)phenyl]-6-(trifluoromethyl) pyrimidine (5.78g) in MeOH (500ml) was added a solution of OXONE™ (Aldrich, 56.23g, 5eq) in water (200ml). The mixture was stirred at ambient temperature overnight, concentrated *in vacuo* and the residue partitioned between water and ethyl acetate (2 x 100ml). The combined organic phases were dried and concentrated *in vacuo* to an off-white solid which was triturated with hot isopropanol to give the title compound as a white solid (5.6g, 80%).

MH+ 381

Tlc SiO<sub>2</sub> Ethyl acetate:cyclohexane (1:1) R<sub>f</sub> 0.45


Example 12-(4-Fluorophenoxy)-4-[4-(methylsulfonyl)phenyl]-6-(trifluoromethyl)pyrimidine.

To a stirred solution of 4-fluorophenol (37mg, 0.33mmole) in dry tetrahydrofuran (10ml) was added, under an atmosphere of nitrogen, sodium hydride (60% dispersion in oil, 13mg, 0.33mmole) and the resulting mixture stirred at 20 for 30min. To the stirred reaction mixture was added 2-(methylsulfonyl)-4[4-(methylsulfonyl)phenyl]-6-trifluoromethyl)pyrimidine (114mg, 0.33mmole) in a single portion, and stirring was continued for 2h. The solvent was evaporated, and the residue partitioned between dichloromethane and 2N sodium hydroxide. The dried organic phase was evaporated to dryness. The residue was purified on a silica gel SPE cartridge eluting with chloroform to afford the title compound as a colourless solid (99mg, 80%).

MH+ 413.


Examples 2 to 10

- Examples 2 to 10, as shown in Table 1 that follows, were prepared in the manner described for Example 1.

**Table 1**

Ex	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	MS
2	3,4-difluorophenyl	CF <sub>3</sub>	CH <sub>3</sub>	MH+ 431
3	4-methoxyphenyl	CF <sub>3</sub>	CH <sub>3</sub>	MH+ 425
4	4-fluorobenzyl	CF <sub>3</sub>	CH <sub>3</sub>	MH+ 427
5	4-bromophenyl	CF <sub>3</sub>	CH <sub>3</sub>	MH+ 474
6	4-methylphenyl	CF <sub>3</sub>	CH <sub>3</sub>	MH+ 409
7	5-chloropyridin-3-yl	CF <sub>3</sub>	CH <sub>3</sub>	MH+ 431
8	cyclohexyl	CF <sub>3</sub>	CH <sub>3</sub>	MH+ 401

Table 1



Ex	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	MS
9	cyclopentylmethyl	CF <sub>3</sub>	CH <sub>3</sub>	MH+ 401
10	n-butyl	CF <sub>3</sub>	CH <sub>3</sub>	MH+ 375

Example 112-Butoxy-4-[4-(methylsulfonyl)phenyl]-6-(trifluoromethyl)pyrimidine

Sodium methoxide (6.6kg of a 30%w/w solution in methanol) was added over at least 30min to a solution of 4-(methylthio)acetophenone (5.0kg) and methyl trifluoroacetate (4.25kg) in tert-butylmethylether (40L) at 40±3°C. The solution was heated at 40±3°C for at least 3h. Acetic acid (55L) was added, followed by S-methyl 2-thiopseudourea sulfate (5.45kg) and the mixture concentrated to ca. 45L. The mixture was heated at about 110°C for at least a further 8h (overnight) then acetic acid (20L) was added before cooling to 50±3°C. A solution of sodium tungstate dihydrate (0.2kg) in water (2.5L) was added, followed by hydrogen peroxide (20.7kg of 30%w/v solution), which was added over at least 3h, maintaining the temp at ca. 50°. The mixture is heated at ca. 50°C for at least 12h before cooling to 20±3°C. A solution of sodium sulphite (3.45kg) in water (28L) was then added over at least 30min whilst maintaining the temperature at 20±3°. The mixture was aged at 20±3°C for ca. 1h and 2-(methylsulfonyl)-4-[4-(methylsulfonyl)phenyl]-6-(trifluoromethyl)pyrimidine collected by filtration, washed with water (3x15L) and dried at up to 60° *in vacuo*. Yield, 9.96kg, 90% of theory.

20

A suspension of 2-(methylsulfonyl)-4-[4-(methylsulfonyl)phenyl]-6-(trifluoromethyl)pyrimidine (525g) in n-butanol (5.25L) was treated with potassium carbonate (210g) at 20±5°C. The mixture was heated to 50±5°C overnight until the reaction was complete by HPLC. Acetic acid (1.57L) was added dropwise, to control any gas evolution, keeping the temperature at

25

50±5°C. Water (3.67L) was then added over 30min keeping the temperature at 50±5°C to allow full crystallisation to occur. The slurry was then cooled to 20-25°C and aged for at least 1 hour. The resulting product was then filtered under vacuum and washed with a mixture of n-butanol (787mL), acetic acid (236mL), and water (551mL) followed by water (2x1.57L). The product was then dried at up to ca50°C under vacuum to yield the title compound. Yield, 457g, 88.4% of theory. The title compound was found to be identical to that of Example 10.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 8.33(2H, d, para-di-substituted CH); 8.11(2H, d, para-di-substituted CH); 7.70(1H, s, aromatic CH); 4.54(2H, t, butyl CH<sub>2</sub>); 3.12(3H, s, sulphone CH<sub>3</sub>); 1.88(2H, m, butyl CH<sub>2</sub>); 1.55(2H, m, butyl CH<sub>2</sub>); 1.01(3H, t, butyl CH<sub>3</sub>).

### Biological Data

#### 15 Cell Based Assay

Inhibitory activity against human COX-1 and COX-2 was assessed in COS cells which had been stably transfected with cDNA for human COX-1 and human COX-2. 24 Hours prior to experiment, COS cells were transferred from the 175cm<sup>2</sup> flasks in which they were grown, onto 24-well cell culture plates using the following procedure. The incubation medium (Dulbecco's modified eagles medium (DMEM) supplemented with heat-inactivated foetal calf serum (10%v/v), penicillin (100 IU/ml), streptomycin (100µg/ml) and geneticin (600µg/ml)) was removed from a flask of confluent cells (1 flask at confluency contains approximately 1x10<sup>7</sup> cells). 5ml of phosphate buffered saline (PBS) was added to the flask to wash the cells. Having discarded the PBS, cells were then incubated with 5ml trypsin for 5 minutes in an incubator (37°). The flask was then removed from the incubator and 5ml of fresh incubation medium was added. The contents of the flask was transferred to a 250ml sterile container and the volume of incubation medium subsequently made up to 100ml. 1ml cell suspension was pipetted into each well of 4x24-well cell culture plates. The plates were then placed in an incubator (37°C, 95% air/5% CO<sub>2</sub>) overnight. If more than 1 flask of cells were required, the cells from the individual flasks were combined before being dispensed into the 24-well plates.

Following the overnight incubation, the incubation medium was completely removed from the 24-well cell culture plates and replaced with 250µl fresh DMEM (37°C). The test compounds were made up to 250x the required test

concentration in DMSO and were added to the wells in a volume of 1 $\mu$ l. Plates were then mixed gently by swirling and then placed in an incubator for 1 hour (37°C, 95% air/5% CO<sub>2</sub>). Following the incubation period, 10 $\mu$ l of arachidonic acid (750 $\mu$ M) was added to each well to give a final arachidonic acid concentration of 30 $\mu$ M. Plates were then incubated for a further 10 minutes, after which the incubation medium was removed from each well of the plates and stored at -20°C, prior to determination of prostaglandin E<sub>2</sub> (PGE<sub>2</sub>) levels using enzyme immunoassay. The inhibitory potency of the test compound was expressed as an IC<sub>50</sub> value, which is defined as the concentration of the compound required to inhibit the PGE<sub>2</sub> release from the cells by 50%. The selectivity ratio of inhibition of COX-1 versus COX-2 was calculated by comparing respective IC<sub>50</sub> values.

The following IC<sub>50</sub> values for inhibition of COX-2 and COX-1 were obtained from the cell based assay for compounds of the invention:

Example No.	COX-2: IC <sub>50</sub> (nM)	COX-1: IC <sub>50</sub> (nM)
1	<1	81,300
2	23	9,675
3	4	2,923
5	6	61,380

#### 15 Microsomal Assay

Inhibitory activity against microsomal h-COX2 was assessed against a microsomal preparation from baculovirus infected SF9 cells. An aliquot of microsomal preparation was thawed slowly on ice and a 1/40,000 dilution prepared from it into the assay buffer (sterile water, degassed with argon containing 100mM HEPES (pH 7.4), 10mM EDTA (pH7.4), 1mM phenol, 1mM reduced glutathione, 20mg/ml gelatin and 0.001mM Hematin). Once diluted the enzyme solution was then sonicated for 5 seconds (Branson sonicator, setting 4, 1cm tip) to ensure a homogeneous suspension. 155 $\mu$ l enzyme solution was then added to each well of a 96-well microtitre plate containing either 5 $\mu$ l test compound (40x required test concentration) or 5 $\mu$ l DMSO for controls. Plates were then mixed and incubated at room temperature for 1 hour. Following the incubation period, 40 $\mu$ l of 0.5 $\mu$ M arachidonic acid was added to each well to give


a final concentration of 0.1 $\mu$ M. Plates were then mixed and incubated for exactly 10 minutes (room temperature) prior to addition of 25 $\mu$ l 1M HCl (hydrochloric acid) to each well to stop the reaction. 25 $\mu$ l of 1M NaOH (sodium hydroxide) was then added to each well to neutralise the solution prior to determination of PGE<sub>2</sub> levels by enzyme immunoassay (EIA).

The following IC<sub>50</sub> values for inhibition of COX-2 and COX-1 were obtained from the microsomal assay for compounds of the invention:

Example No.	COX-2: IC <sub>50</sub> (nM)	COX-1: IC <sub>50</sub> (nM)
6	<10	3,752
7	<10	79,889
8	<10	1,860
9	22	69,000
10	22	>30000

CLAIMS

## 1. Compounds of formula (I)



5 in which:

$R^1$  is selected from the group consisting of H,  $C_{1-6}$ alkyl,  $C_{1-2}$ alkyl substituted by one to five fluorine atoms,  $C_{3-6}$ alkenyl,  $C_{3-6}$ alkynyl,  $C_{3-10}$ cycloalkyl,  $C_{0-6}$ alkyl,  $C_{4-12}$ bridged cycloalkyl,  $A(CR^4R^5)_n$  and  $B(CR^4R^5)_n$ ;

$R^2$  is  $C_{1-2}$ alkyl substituted by one to five fluorine atoms;


10  $R^3$  is selected from the group consisting of  $C_{1-6}$ alkyl,  $NH_2$  and  $R^7CONH$ ;

$R^4$  and  $R^5$  are independently selected from H or  $C_{1-6}$ alkyl;

A is an unsubstituted 5- or 6-membered heteroaryl or an unsubstituted 6-membered aryl, or a 5- or 6-membered heteroaryl or a 6-membered aryl substituted by one or more  $R^6$ ;

15  $R^6$  is selected from the group consisting of halogen,  $C_{1-6}$ alkyl,  $C_{1-6}$ alkyl substituted by one more fluorine atoms,  $C_{1-6}$ alkoxy,  $C_{1-6}$ alkoxy substituted by one or more F,  $NH_2SO_2$  and  $C_{1-6}alkylSO_2$ ;

B is selected from the group consisting of



) defines the point of attachment of the ring;

20


$R^7$  is selected from the group consisting of H,  $C_{1-6}$ alkyl,  $C_{1-6}$ alkoxy,  $C_{1-6}alkylOC_{1-6}alkyl$ , phenyl,  $HO_2CC_{1-6}alkyl$ ,  $C_{1-6}alkylOCOC_{1-6}alkyl$ ,  $C_{1-6}alkylOCO$ ,  $H_2NC_{1-6}alkyl$ ,  $C_{1-6}alkylOCONHC_{1-6}alkyl$  and  $C_{1-6}alkylCONHC_{1-6}alkyl$ ; and

25

n is 0 to 4.

2. Compounds as claimed in claim 1 wherein  $R^1$  is selected from the group consisting of  $C_{1-6}$ alkyl,  $C_{1-2}$ alkyl substituted by one to five fluorine atoms,  $C_{3-10}$ cycloalkyl $C_{0-6}$ alkyl and  $A(CR^4R^5)_n$ .
- 5 3. Compounds as claimed in claim 1 or 2 wherein  $R^2$  is  $CHF_2$ ,  $CH_2F$  or  $CF_3$ .
4. Compounds as claimed in any of claims 1 to 3 wherein  $R^3$  is  $C_{1-6}$ alkyl.
5. Compounds as claimed in any of claims 1 to 4 wherein  $R^4$  and  $R^5$  are independently selected from H or methyl.
6. Compounds as claimed in any of claims 1 to 5 wherein A is selected from the group consisting of

10




) defines the point of attachment of the ring  
and A is unsubstituted or substituted by one or two  $R^6$ .

7. Compounds as claimed in any of claims 1 to 6 wherein  $R^6$  is selected from the group consisting of halogen,  $C_{1-3}$ alkyl,  $C_{1-3}$ alkyl substituted by one to three fluorine atoms, and  $C_{1-3}$ alkoxy.
- 15 8. Compounds as claimed in any of claims 1 to 7 wherein  $R^7$  is selected from the group consisting of  $C_{1-6}$ alkyl, phenyl and aminomethyl.
9. Compounds as claimed in any of claims 1 to 8 wherein n is 0 to 2.
10. Compounds as claimed in any of claims 1 to 9 wherein  $R^1$  is  $C_{1-6}$ alkyl;  $R^2$  is  $CF_3$ ; and  $R^3$  is  $C_{1-6}$ alkyl, such as  $C_{1-3}$ alkyl.
- 20 11. Compounds as claimed in any of claims 1 to 10 wherein  $R^1$  is  $C_{3-10}$ cycloalkyl $C_{0-6}$ alkyl, such as  $C_{3-10}$ cycloalkyl;  $R^2$  is  $CF_3$ ; and  $R^3$  is  $C_{1-6}$ alkyl, such as  $C_{1-3}$ alkyl.

12. Compounds as claimed in any of claims 1 to 11 wherein  $R^1$  is  $C_{3-10}$ cycloalkylmethyl, such as  $C_{3-7}$ cycloalkylmethyl;  $R^2$  is  $CF_3$ ; and  $R^3$  is  $C_{1-6}$ alkyl, such as  $C_{1-3}$ alkyl.

5 13. Compounds as claimed in any of claims 1 to 12 wherein  $R^1$  is  $A(CR^4R^5)_n$ ;  $R^2$  is  $CF_3$ ;  $R^3$  is methyl;  $R^4$  and  $R^5$  are both H; A is selected from the group consisting of




10 and A is unsubstituted or substituted by one or two  $R^6$ ;  $R^6$  is selected from the group consisting of fluorine, chlorine, methyl,  $CF_3$  and methoxy; and n is 0 or 1.

14. Compounds of formula (I) as described in Examples 1 to 10.

15. 2-Butoxy-4-[4-(methylsulfonyl)phenyl]-6-(trifluoromethyl)pyrimidine.

15 16. A process for the preparation of a compound of formula (I) as defined in any one of claims 1 to 15, which comprises:

(A), reacting an alcohol  $R^1OH$  of formula (II) or a protected derivative thereof with a compound of formula (III)



20 and thereafter and if necessary,

(B), interconverting a compound of formula (I) into another compound of formula (I); and/or

(C), deprotecting a protected derivative of compound of formula (I).

17. A pharmaceutical composition comprising a compound of formula (I) as defined in any one of claims 1 to 15 in admixture with one or more physiologically acceptable carriers or excipients.
- 5 18. A compound of formula (I) as defined in any one of claims 1 to 15 for use in human or veterinary medicine.
- 10 19. A method of treating a human or animal subject suffering from a condition which is mediated by COX-2 which comprises administering to said subject an effective amount of a compound of formula (I) as defined in any one of claims 1 to 15.
20. A method of treating a human or animal subject suffering from an inflammatory disorder, which method comprises administering to said subject an effective amount of a compound of formula (I) as defined in any one of claims 1 to 15.
- 15 21. The use of a compound of formula (I) as defined in any one of claims 1 to 15 for the manufacture of a therapeutic agent for the treatment of a condition which is mediated by COX-2.
- 20 22. The use of a compound of formula (I) as defined in any one of claims 1 to 15 for the manufacture of a therapeutic agent for the treatment of an inflammatory disorder.

## INTERNATIONAL SEARCH REPORT

Intd      nal Application No

PCT/GB 02/02415

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7    C07D239/34    C07D401/12    A61K31/505    A61P29/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7    C07D    A61K    A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 98 03484 A (GAUTHIER JACQUES YVES ;MERCK FROSST CANADA INC (CA); DUBE DANIEL ( ) 29 January 1998 (1998-01-29) examples, abstract, claims tables 1,2 ---	1-22
Y	US 3 592 895 A (HEPWORTH WALTER ET AL) 13 July 1971 (1971-07-13) claims ---	1-22
P,A	WO 01 38311 A (HARTLEY CHARLES DAVID ;PAYNE JEREMY JOHN (GB); PEGG NEIL ANTHONY ( ) 31 May 2001 (2001-05-31) the whole document --- -/-	1-22

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents :

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*&amp;\* document member of the same patent family

Date of the actual completion of the international search

18 September 2002

Date of mailing of the international search report

26/09/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Stroeter, T

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 02/02415

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, A	WO 01 58881 A (PAYNE JEREMY JOHN ; PEGG NEIL ANTHONY (GB); NAYLOR ALAN (GB); GLAXO) 16 August 2001 (2001-08-16) the whole document -----	1-22

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/GB 02/02415

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
  
Although claims 19, 20 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 02/02415

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9803484	A	29-01-1998	AU 723179 B2	17-08-2000
			AU 3331997 A	10-02-1998
			BG 103179 A	30-11-1999
			CA 2260016 A1	29-01-1998
			WO 9803484 A1	29-01-1998
			CZ 9900130 A3	16-06-1999
			EE 9900018 A	16-08-1999
			EP 0912518 A1	06-05-1999
			HR 970389 A1	30-06-1998
			JP 11514008 T	30-11-1999
			JP 3251945 B2	28-01-2002
			JP 2002080453 A	19-03-2002
			NO 990191 A	16-03-1999
			NZ 333230 A	25-08-2000
			PL 330995 A1	21-06-1999
			SK 3699 A3	10-12-1999
			TR 9900046 T2	21-04-1999
			TW 453994 B	11-09-2001
			BR 9710372 A	17-08-1999
			HU 9903974 A2	28-03-2000
			US 6001843 A	14-12-1999
			US 6071936 A	06-06-2000
			US 5861419 A	19-01-1999
US 3592895	A	13-07-1971	GB 1121922 A	31-07-1968
			BE 700109 A	18-12-1967
			DE 1774407 A1	15-07-1971
			DK 115917 B	24-11-1969
			DK 117836 B	08-06-1970
			ES 341927 A1	01-07-1968
			FR 7509 M	15-12-1969
			FR 1571589 A	20-06-1969
			GB 1157157 A	02-07-1969
			NL 6708449 A	18-12-1967
			US 3502673 A	24-03-1970
			US 3554421 A	12-01-1971
			DK 119089 B	09-11-1970
			DK 118560 B	07-09-1970
			DK 116217 B	22-12-1969
			DK 116218 B	22-12-1969
WO 0138311	A	31-05-2001	AU 2508401 A	04-06-2001
			BR 0015821 A	30-07-2002
			WO 0138311 A2	31-05-2001
			EP 1235812 A2	04-09-2002
			NO 20022470 A	18-07-2002
WO 0158881	A	16-08-2001	AU 3203601 A	20-08-2001
			WO 0158881 A1	16-08-2001